

Name	Other names	Frequency	When can treatment be started?	Pharmacy prior auth. needed?
Methadone	Dolophine [®] , Methadose [®] (oral)	Daily	As soon as indicated, however initial dose should be low and carefully monitored to avoid sedation and/or overdose	N/A
Buprenorphine	Suboxone [®] (buprenorphine & naloxone film)	Daily	As soon as indicated, however should await mild-moderate withdrawal from opioids to avoid precipitated withdrawal	No
	Generic buprenorphine & naloxone (tablet & film), Bunavail [®] (buprenorphine & naloxone film), and Zubsolv [®] (buprenorphine & naloxone tablet)	Daily		Yes
	Sublocade [®] (extended-release buprenorphine injection)	Monthly	After being on oral buprenorphine product for at least seven days	Yes
	Probuphine [®] (buprenorphine implant)	Every six months	After being on oral buprenorphine product for at least three months	Yes
	Buprenorphine monoprodukt (previously branded Subutex [®]) (tablet)	Daily	As soon as indicated, however should await mild-moderate withdrawal from opioids to avoid precipitated withdrawal	No
Naltrexone	Naltrexone (previously branded Revia [®]) (tablet)	Daily	Minimum of seven to 10 days since last opioid use (10-14 days if previously on methadone or buprenorphine), otherwise will trigger opioid withdrawal	No
	Vivitrol [®] (extended-release naltrexone injection)	Monthly		No